

JOURNAL OF PLASMA AND FUSION RESEARCH

The Journal of the Japan Society of Plasma Science and Nuclear Fusion Research
Vol. 86, No.2, February 2010

Special Topic Article

Recent Progress of Ion Heat Transport Study on LHD

| | | |
|---|--|----|
| 1. Introduction | NAGAOKA Kenichi | 69 |
| 2. Improvement of Ion Heat Transport on LHD | NAGAOKA Kenichi and YOKOYAMA Masayuki | 71 |
| 3. Observations of Spontaneous Toroidal Flow and Impurity Hole on LHD | YOSHINUMA Mikirou | 78 |
| 4. Comparison with an Internal Transport Barrier (ITB) in Tokamaks | IDA Katsumi | 86 |
| 5. Prospects on High-Ion-Temperature Plasmas in the LHD | YOKOYAMA Masayuki and TAKEIRI Masahiko | 94 |

Lecture Note

R&D Activities for 30 Years on Handling Technology of a Large Amount of Tritium and Future Subjects

| | | |
|---|---|----|
| 1. Establishment of Facility for Handling a Large Amount of Tritium – The Dawn of Tritium Studies in Japan and Towards the Facility Construction – | MATSUYAMA Masao and YAMANISHI Toshihiko | 97 |
|---|---|----|

Research and Technology Note

| | | |
|---|-----------------|-----|
| Development of High-Power Microwave Tubes for Fusion Experimental Devices | HAYASHI Kenichi | 104 |
|---|-----------------|-----|

| | | |
|----------------------------|--|-----|
| PFR Abstracts | | 122 |
|----------------------------|--|-----|

| | | |
|--------------------------|--|-----|
| Information | | 123 |
|--------------------------|--|-----|

| | | |
|---|--|-----|
| Plasma & Fusion Calendar | | 132 |
|---|--|-----|

| | | |
|---------------------------|--|-----|
| Announcement | | 134 |
|---------------------------|--|-----|

Cover

SEM pictures of carbon dust observed on the graphite target irradiated by Ar/H₂/N₂ plasmas at different N₂ injection ratio to hydrogen. The injection of a small amount of nitrogen gas led to significant suppression of carbon dust formation on the graphite surface. With increasing N₂ injection ratio, the carbon dust shape changes into polyhedral particles at an N₂ content of 0.3 – 0.7% and clusters made of smaller particles at an N₂ content > ~ 2 %. (Masaaki KYO *et al.*, Plasma and Fusion Research Vol.5, 004 (2010) <http://www.jspf.or.jp/PFR/>)

Published Monthly by

The Japan Society of Plasma Science and Nuclear Fusion Research

3-1-1, Uchiyama, Chikusa-ku, Nagoya 464-0075, Japan

Tel 052-735-3185, Fax 052-735-3485, E-mail: plasma@jspf.or.jp, URL: <http://www.jspf.or.jp/>