

JOURNAL OF PLASMA AND FUSION RESEARCH

The Journal of the Japan Society of Plasma Science and Nuclear Fusion Research

Vol. 89, No.5, May 2013

Front Runner

Spectroscopy and Atomic Structure of High-Z Multiply-Charged Ions in High-Temperature Plasmas	
1. Introduction	MORITA Shigeru 271
2. Spectroscopic Observation of Multiply Charged Tungsten Ions in Magnetically Confined Plasmas	HASUO Masahiro, TANAKA Hirotaka and YANAGIBAYASHI Jun 273
3. EUV Spectroscopy of High-Z Multiply Charged Ions in LHD	SUZUKI Chihiro 284
4. Study of Tungsten Spectra in LHD	MORITA Shigeru, SAKAUE Hiroyuki, DONG Chunfeng, GOTO Motoshi and OISHI Tetsutarou 289
5. Observation of Visible Forbidden Lines from High-Z Multiply Charged Ions at LHD	KATO Daiji 296
6. Production and Spectroscopy of High-Z Multiply Charged Ions with EBIT	NAKAMURA Nobuyuki and SAKAUE Hiroyuki 299
7. Theory of Atomic Structure for High-Z Multiply Charged Ions	KOIKE Fumihiko 305
8. Present Status of the Modeling of High-Z Multiply Charged Ions	SASAKI Akira 308
9. Spectroscopic Model for Tungsten Ions	MURAKAMI Izumi 314
10. Concluding Remarks	MURAKAMI Izumi 318

Commentary

Plasma Boundary of 3D MHD Equilibrium Inferred from Flow Patterns	IDA Katsumi, KAMIYA Kensaku and SUZUKI Yasuhiro 319
--	---

Information	328
-------------------	-----

Plasma & Fusion Calendar	332
--------------------------------	-----

Announcement	333
--------------------	-----

Cover

Target injection system for laser fusion. A target loader stores more than 10,000 CD targets. The target inside the target loader falls onto a rotating disk and is conveyed to the exit hole and flies 18 cm down to the laser focal point. Flying target is engaged by laser beams at 1 Hz. (Osamu KOMEDA *et al.*, Plasma and Fusion Research Vol.8, 1205020 (2013) <http://www.jspf.or.jp/PFR/>)