

JOURNAL OF PLASMA AND FUSION RESEARCH

The Journal of the Japan Society of Plasma Science and Nuclear Fusion Research
Vol. 94, No.7, July 2018

Commentary

- Simulated Annealing for MHD Equilibria with Flow and/or Magnetic Island FURUKAWA Masaru 341

Lecture Note

- Liquid Metal as the Neutron Source
1. Introduction TAKAHASHI Mitsutoshi 349
2. The Handling of the Liquid Metals for Neutron Sources MIURA Kuniaki 350

PFR Abstracts

- Mourning 355

- Information 359

- Plasma & Fusion Calendar 367

- Announcement 368

Cover

Snap shots of nanoparticle transport between a plasma source and a substrate holder for (a) gas velocity of 0.18 m/s and (b) that of 0.73 m/s. For 0.18 m/s, nanoparticles are trapped between plasma generated in a multi-hollow electrode and a substrate at 2 s after discharge initiation. Then, they are transported toward the substrate after 50 s. The trapped nanoparticles might play as scavengers which block the transport of nanoparticles. For 0.73 m/s, nanoparticles are transported just after the plasma ignition.(Takashi KOJIMA *et al.*, Plasma and Fusion Research, Vol.13, 1406082 (2018) <http://www.jspf.or.jp/>)

Published Monthly by

The Japan Society of Plasma Science and Nuclear Fusion Research

3-1-1 Uchiyama, Chikusa-ku, Nagoya 464-0075, Japan

Tel (052)735-3185, Fax (052)735-3485, E-mail: plasma@jspf.or.jp, URL: <http://www.jspf.or.jp/>