# Chromatographic Analysis of Molecular Hydrogen (H<sub>2</sub>) in the Atmosphere for Understanding Atmospheric Tritiated Hydrogen (HT)\*)

### Masahiro TANAKA, Chie IWATA, Miki NAKADA, Saori KURITA and Hideki KAKIUCHI<sup>1)</sup>

National Institute for Fusion Science, National Institutes of Natural Sciences, Oroshi, Toki, Gifu 509-5292, Japan <sup>1)</sup>Institute for Environmental Sciences, Rokkasho, Kamikita, Aomori 039-3212, Japan (Received 22 December 2022 / Accepted 10 April 2023)

Tritium monitoring would be required at the site of the fusion test facility from the viewpoints of radiation safety and public acceptance. As one of the tritium monitoring items, atmospheric tritium monitoring in the environment is conducted and has three chemical forms: tritiated water vapor (HTO), tritiated molecular hydrogen (HT), and tritiated methane (CH<sub>3</sub>T). It is well-known that the specific activity of HT and CH<sub>3</sub>T is much higher than that of HTO. To investigate the cause of high specific activity, we focus on the behavior of molecular hydrogen (H<sub>2</sub>) in the atmosphere and develop a measurement system based on a gas chromatograph. Monitoring results showed that the range of the H<sub>2</sub> mixing ratio at the NIFS Toki site was  $0.4 \sim 0.6$  ppm over the observation period. Atmospheric H<sub>2</sub> concentrations were higher during the daytime and long daylight seasons, suggesting the generation of hydrogen by photochemical reactions. Preliminary results suggest that there does not appear to be a clear correlation between the concentration of atmospheric HT and the concentration of H<sub>2</sub> in the atmosphere. This suggests that the source of HT is different from the source of H<sub>2</sub> in the atmosphere.

© 2023 The Japan Society of Plasma Science and Nuclear Fusion Research

Keywords: trace constituents of the atmosphere, atmospheric tritium, atmospheric molecular hydrogen, gas chromatograph system, trace reduction detector, hydrogen flame ionization detector

DOI: 10.1585/pfr.18.2405038

## 1. Introduction

Tritium, which is a radioactive hydrogen isotope, is a cosmogenic nuclide produced by the nuclear reactions between the atmosphere (N<sub>2</sub>, O<sub>2</sub>) and cosmic rays. The amount of tritium in nature is estimated to be approximately 1-1.3 EBq. On the other hand, the artificial tritium increased due to the atmospheric nuclear bomb testing from 1945 to 1980, and the artificial tritium amount was estimated to be approximately 186 EBq [1]. In recent years, the impact of tritium released from atomic power stations such as heavy water reactor and fuel reprocessing facilities was suggested [1, 2]. In addition, nuclear fusion as the prospective new energy source will become a large tritium release source into the environment.

To understand the plasma physics and progress the fusion reactor development, deuterium gas is utilized as operational gas in the large fusion test device such as Large Helical Device (LHD) in National Institute for Fusion Science (NIFS) and then a small amount of tritium is produced by D-D fusion reaction in the vacuum vessel [3]. Thus, tritium exhausted from LHD would be recovered by an exhaust detritiation system (EDS) [4]. On the other hand, environmental tritium monitoring at the NIFS Toki site is required from the viewpoint of public acceptance. At the NIFS Toki site, tritium in river water, pond water, well water, tap water, rainwater, pine needle, and atmosphere has been measured for a few decades [5-7]. As for atmospheric tritium, the chemical forms of tritium are water vapor (HTO), molecular hydrogen (HT), and hydrocarbons (mainly CH<sub>3</sub>T) [8–10]. Thus, the atmospheric tritium monitoring system with discriminating chemical forms was developed and the variation of atmospheric tritium at the NIFS Toki site has been observed since 2004 [11]. It is known that the specific activities of HT and CH<sub>3</sub>T are orders of magnitude greater than that of HTO. To investigate the cause of high specific activity, we focus on the behavior of molecular hydrogen  $(H_2)$  in the atmosphere. Recently, atmospheric H<sub>2</sub> has attracted interest in association with greenhouse gas and from the viewpoint of the creation of a hydrogen energy society. The mixing ratio of  $H_2$  in the atmosphere is reported to be ~0.53 ppm and the lifetime is estimated to be about 2 years. The sources of H<sub>2</sub> are mainly fossil fuel combustion, biomass burning, and the photochemical degradation of organic compounds within the troposphere. The sinks of  $H_2$  are soil uptake by enzymes and microorganisms, and the reaction of H<sub>2</sub> with the OH radical in the troposphere [12-14]. Assuming that the variation of HT is correlated with that of H<sub>2</sub> in the atmosphere, the source of HT would be the same as that of H<sub>2</sub>. In this report, we developed the monitoring system of

author's e-mail: tanaka.masahiro@nifs.ac.jp

<sup>&</sup>lt;sup>\*)</sup> This article is based on the presentation at the 31st International Toki Conference on Plasma and Fusion Research (ITC31).

trace constituents of the atmosphere,  $H_2$ , CO, and  $CH_4$ , and then measured them with atmospheric tritium monitoring for a few months.

#### 2. Monitoring System

The development of a gas chromatograph system for continuous trace analysis of H<sub>2</sub>, CO, and CH<sub>4</sub> in the atmosphere was based on the reports by Kawano et al., [15, 16]. The flow diagram of the gas chromatography system is shown in Fig. 1. The system consists of a gas chromatograph (G2700TF with hydrogen flame ionization detector (FID) for the detection of CH<sub>4</sub>, GTR TEC Corporation), an autosampler with a sampling pump and two multiport valves, and trace reduction detector for the detection of H<sub>2</sub> and CO (TRD, J-SCIENCE LAB Co., Ltd.). The sampling control and chromatogram analysis were conducted by HCP software (Version: 5.0.2.354, GTR TEC corporation). The gas separation column was a series of molecular sieves 5A (MS-5A, column length: 2m) and Porapak-Q (PQ, column length: 2 m). The carrier gas was ultra-high purity nitrogen gas (Purity, >99.99995%). The ultra-high purity hydrogen gas (Purity, >99.99999%) and purified air by passing through a catalyst heated up to 500 °C to burn off all hydrocarbons in the atmosphere were used for FID. The sampling gas volume was  $2 \text{ cm}^3$ . The detection ranges of FID and TRD were more than sub-ppm order for CH<sub>4</sub> and a few ten ppb to 10 ppm for  $H_2$  and CO. The measurement interval was 2 or 3 hr. The height of the monitoring point was 6 m from ground level. To detect the different gas species by two detectors and discriminate oxygen component, 6 ports valve was switched on the way of chromatographic measurement. An example of a chromatogram is shown in Fig. 2. The first large peak is hydrogen, and the second and third large peaks are methane and carbon monoxide. The chromatogram indicates sufficient retention time for the discrimination of hydrogen, methane, and carbon monoxide. The standard gas of 0.98 ppm of H<sub>2</sub>, CO, and 2.0 ppm of CH<sub>4</sub> based on N<sub>2</sub> was used for the calibration of the chromatogram. The calibration operation was conducted once a month and the measurement results were corrected. The gas sampling line of the gas chromatograph system was connected to the sampling line of the original tritium active sampler system. The detailed specifications, operation conditions of the tritium active sampler system, and tritium measurement are described in the reports of Uda *et al.* and Tanaka *et al.* [11, 17].

The meteorological observation station was installed on the top of the research staff building at the center of the Toki site. The height of the observation point was approximately 40 m from the ground level. The observation items were wind direction, wind speed, atmospheric pressure, air temperature, humidity, ground temperature, solar radiation, and precipitation. The data acquisition interval



Fig. 2 An example of chromatogram by the gas chromatography using FID and TRD.



Fig. 1 Schematic diagram of gas flow line in the gas chromatograph and the autosampler.

was set to be 10 seconds.

The NIFS Toki site is located about 100 km inland, north of the Pacific coastal areas, and the buildings stand on a hill at an altitude of about 200 m. The measurement of trace constituents of the atmosphere was started concurrently with atmospheric tritium monitoring in late June 2022.

## 3. Results and Discussion

The concentration variation of the trace constitutions of the atmosphere for four months is shown in Fig. 3. The measurement was interrupted for the separation column conditioning from 4th October to 7th October. Large peak concentration was occasionally observed in all trace constitutions. Diurnal and long-term variation of  $H_2$  concentration was observed. The concentrations of CO and  $CH_4$ were not indicated long-term variation and were almost constant.

To clarify the diurnal variations, the concentration at each measurement time in a day was averaged from June to October, and the results are shown in Fig. 4. Here, the time average and integrated solar radiation in a day were also indicated. H<sub>2</sub> concentration was the lowest at 6:00 and then gradually increases during the daytime, and then reached a maximum around 18:00~20:00 at sunset. The solar radiation began the increase at 6:00 and reached a peak at noon. After 18:00, the solar radiation became almost zero. The variation of H<sub>2</sub> concentration seems to correlate with the integrated solar radiation. During the daytime, photochemical reactions are supposed to promote the production of H<sub>2</sub>. Then, H<sub>2</sub> uptake in the soil would occur in the nighttime and the H<sub>2</sub> concentration might decrease. On the other hand, CH<sub>4</sub> concentration slightly increased during the nighttime and decreased during the daytime. The variation in CO concentration was similar to that of H<sub>2</sub>. They suppose that the correlation between CH<sub>4</sub> and H<sub>2</sub> has negative and the correlation between CO and H<sub>2</sub> has positive. These correlations indicate that the photochemical reaction to produce H<sub>2</sub> in the atmosphere starts from CH<sub>4</sub> or VOC, and then some of the final products are CO and H<sub>2</sub> [14].

Figure 5 shows the variations of monthly mean solar radiation, the concentration of trace constitutions in the atmosphere, and HT concentration from June to November. The shorter the daytime toward winter, the lesser the solar radiation.  $H_2$  concentrations reached a maximum in July, and a gradual decrease was observed through the fall. Except for June data,  $H_2$  concentration would be correlated with solar radiation. However, the correlation between  $H_2$  and other species of CO and CH<sub>4</sub> observed in diurnal variation data was not obtained clearly in the monthly mean data. A part of the concentration variation would be caused by the photochemical reactions due to solar radiation as well as the temporary effect due to air mass such as migratory anticyclones and cyclones from the Asian Continent



Fig. 3 Concentration variations of trace constitutions of the atmosphere at Toki: (a) Hydrogen, (b) Carbon monoxide, (c) Methane.



Fig. 4 Average diurnal variations of solar radiation and the concentration of trace constitutions of the atmosphere.

and the Pacific Ocean. The air mass transports an atmosphere from another area and mixes with the atmosphere in the observation area. Thus, the seasonal correlation among the species of  $H_2$ , CO, and CH<sub>4</sub> may not be observed.

The concentration of atmospheric HT was almost constant during the observation period. Although these data show preliminary results, it seems that the behavior of atmospheric HT concentration is a different trend from that of atmospheric H<sub>2</sub> concentration. To clarify the correlation between HT and H<sub>2</sub> in the atmosphere, long-term monitor-



Fig. 5 The variations of monthly mean solar radiation, trace constitutions in the atmosphere and molecular tritium (HT): (a) Solar radiation, (b) Concentrations of trace constitution in the atmosphere, (c) Atmospheric HT concentration.

ing of more than one year is required.

## 4. Conclusions

To understand the behavior of atmospheric molecular tritium (HT), a chromatographic analysis of trace constituents (H<sub>2</sub>, CH<sub>4</sub>, CO) of the atmosphere was conducted using a gas chromatograph with a trace reduction detector (TRD) and hydrogen flame ionization detector (FID) at Toki, Japan. H<sub>2</sub> concentrations in the atmosphere were higher during the daytime and the season with long daylight hours, suggesting hydrogen production by photochemical reactions.

As the preliminary result, there does not appear to be a clear correlation between atmospheric HT concentrations and atmospheric  $H_2$  concentrations. It suggests that the source of molecular tritium is different from that of the molecular hydrogen in the atmosphere. For future studies, long-term observational data are needed to obtain clear conclusions.

## Acknowledgments

This study was supported by ERAN (Environmental Radioactivity Research Network Center, University of Tsukuba), Grant Number: F-22-74, and partly JSPS KAK-ENHI, Grant Number: JP21H03607.

- UNSCEAR 2016 Report, Sources, effects and risks of ionizing radiation: Report to the General Assembly, Scientific Annex C, United Nations, New York (2018).
- [2] J.D. Happell et al., Tellus 56B, 183 (2004).
- [3] M. Osakabe et al., Nucl. Fusion 62, 042019 (2022).
- [4] M. Tanaka et al., Fusion Eng. Des. 127, 275 (2018).
- [5] N. Akata et al., Radiat. Prot. Dosim. 167, 210 (2015).
- [6] M. Tanaka and T. Uda, Radiat. Prot. Dosim. 167, 187 (2015).
- [7] N. Akata et al., Plasma Fusion Res. 11, 1305032 (2016).
- [8] T. Okai et al., J. Radioanal. Nucl. Chem. 239, 527 (1999).
- [9] N. Momoshima *et al.*, J. Nucl. Radiochem. Sci. 8, 117 (2007).
- [10] N. Akata et al., J. Env. Radioact. 102, 837 (2011).
- [11] T. Uda et al., Fusion Eng. Des. 81, 1385 (2006).
- [12] P.C. Novelli *et al.*, J. Geophys. Res. **104**, D23, 30427 (1999).
- [13] H. Price et al., J. Geophys. Res. 112, D22, 108 (2007).
- [14] D.H. Ehhalt and F. Rohrer, Tellus **61B**, 500 (2009).
- [15] T. Kawano *et al.*, Jpn. J. Appl. Phys., Part 2: Lett. **42**, L549 (2003).
- [16] T. Kawano et al., J. Chromatogr. A 1023, 123 (2004).
- [17] M. Tanaka et al., Radiat. Prot. Dosim. 198, 1084 (2022).