

ヘリウム混合照射におけるW中のトリチウムリテンションモデリング Modeling tritium retention in tungsten under mixed helium irradiation

遠藤博史, Heun Tae Lee, 大塚裕介, 上田良夫
Hiroshi Endo, Heun Tae Lee, Yusuke Ohtsuka, Yoshio Ueda

大阪大学大学院工学研究科 〒565-0871 大阪府吹田市山田丘 2-1
Graduate School of Engineering Osaka University, 2-1 Yamadaoka, Suita, Osaka, 565-0871, Japan

1. Introduction

Tungsten (W) is a plasma facing material in ITER divertor and candidate armor material for DEMO. The evaluation of tritium (T) retention is important due to its impact on safety and operation. T retention in ITER has been estimated by the previous studies. In the case of D-T phase, neutrons and He ash will be present. On the one hand, neutrons create additional trap sites resulting in an increase in T retention. On the other hand, He can strongly reduce hydrogen retention in W [1] by limiting T diffusion.

In this paper, T retention in W under mixed He irradiation is modeled using TMAP7. In contrast to previous studies, simulations were performed using hydrogen transport parameters benchmarked from permeation experiments [2]. First, the recombination coefficient was determined from D-only irradiation case. Next, recombination coefficient and diffusivity required for mixed D-He irradiation case was fitted. Finally, the benchmarked transport parameters and trap parameters from recent neutron damage study [3] were used to estimate T inventory in ITER-like conditions.

2. Procedure

The detail of the benchmarked experiments is shown. Specimens used in the permeation experiment were polycrystalline W with thickness of 30 μm supplied by A.L.M.T. Corp., Japan. D ion irradiation (D-only case) as well as mixed D-He ion irradiation (D-He case) at 1 keV was performed at $550 \text{ K} \leq T \leq 1050 \text{ K}$. The incident flux varied from 10^{19} - $10^{20} \text{ m}^{-2}\text{s}^{-1}$. The He concentration was fixed at 2 %.

To model the experimental data, the parameters were determined as below. Diffusivity, D , and solubility, S , were taken from Frauenfelder [4]. Initially, Anderl's [5] recombination coefficient was used. However, to fit the experimental data for D-only case, the front segment recombination coefficient, K_1 , needed to be varied. Modification of hydrogen transport during mixed He-D irradiation was modeled by varying the front segment recombination coefficient, K_2 , and diffusivity, D_2 .

Finally, Simulation of ITER-like conditions was done by assuming full W divertor and first walls.

3. Result

Tritium retention with and without He irradiation under ITER-like condition was estimated using TMAP7 code. The surface temperature of W divertor and W first wall were taken as 500 K and 400 K, respectively [6]. The incident flux of W divertor and W first wall were set to be $2.0 \times 10^{23} \text{ m}^{-2}\text{s}^{-1}$ and $1.4 \times 10^{19} \text{ m}^{-2}\text{s}^{-1}$, respectively. The hydrogen transport parameters were benchmarked against ion driven permeation experiments.

Fig. 5 shows the calculated T retention with and without He irradiation. Without He irradiation, the T retention estimated in this work was lower by approximately two orders of magnitude than previous estimate by Roth et al. [6]. This was due to different trap parameters resulting in different trap filling dynamics and due to traps filled sequentially (strong trapping) using 1.8 eV traps. With He irradiation, x 5 reduction in T retention observed compared to D-only case. The calculations were in good agreement with a previous estimate by Wampler et al. [7]




Fig.1 Estimate of T retention in ITER (All W case). The solid lines indicate T retention with and without He irradiation. Plotted in dashed line are previous studies by Roth et al [6] and Wampler and Doerner [7].

Reference

- [1] V. Kh. Alimov et al., J. Nucl. Mater. **420** (2012) 370-373.
- [2] H.T. Lee et al., J. Nucl. Mater. **415** (2011) S696-S700.
- [3] Yuji Hatano et al., 20th Int. Conf. PSI (2012).
- [4] R. Frauenfelder, J. Vac. Sci. Technol. **6** (1969) 388.
- [5] R.A. Anderl et al., Fusion Tech. **21** (1992) 745.
- [6] Roth J. et al., J. Nucl. Mater. **390-391** (2009) 1-9.
- [7] Wampler W R and Doerner R, Nucl. Fusion **49** (2009) 115023.